


1825 Logan Ave. (Entrance 8)
Waterloo, Iowa 50703
319.235.3823

125 E. Tower Park Drive
Waterloo, Iowa 50701
319.234.5990



Ravi Mallavarapu, MD
Srinivas Kalala, MD
Barbara Burkle, ARNP
Tracy Elliott, ARNP
Kelli DeSerano, ARNP

Flatulence (GAS)

What is flatulence?

Flatulence is the passage of air or gas from the intestines through the rectum. The air or gas is called flatus.

It is normal to have some flatulence. The average person passes gas through the rectum 8-20 times a day. Flatulence is usually a problem only when the gas smells foul, happens frequently, or occurs at inconvenient times.

How does it occur?

Part of the flatus in the intestine is air that you have swallowed. You may swallow large amounts of air with your food, especially if you eat or drink quickly. Some people have a nervous habit of swallowing air all day, especially in times of stress. If you are in an upright position, swallowed air can pass back up from your stomach and be released through your mouth in a belch. However, each time you belch, you swallow more air, so the belching is likely to continue. When you lie down, the air may instead pass on through the intestines and rectum and out the anus.

The other cause of gas in the intestines is the digestive process itself. Bacteria in the intestines produce gas as they break down food residues. Some foods are more likely to produce intestinal gas than others, and some medications that cause constipation can make intestinal gas more frequent or painful. Gas formed by bacteria in the digestive process is always passed through the rectum.

What are the symptoms?

The symptoms of flatulence are:

- Frequent expulsion of gas through the anus.
- Abdominal discomfort that is relieved by expulsion of gas.

How is it treated?

Flatulence can be treated with the following measures:

- Try to eat and drink more slowly so you can swallow less air with your food.
- Change your diet. Avoid foods that are known to cause excessive gas.

How can I take care of myself?

You should first realize that some flatulence is normal. Try not to let it bother you. The more you worry about it, the more air you are likely to swallow.

Recognizing that you are swallowing air is the first step in controlling the habit of swallowing air. Chew your food longer and sip fluids slowly. Try to avoid frequent sighing and chewing gum.

A change in diet often can help you have less intestinal gas. Try to eat a more varied diet and avoid foods that cause flatulence. These foods vary from person to person, but the following foods commonly produce gas in many people:

- Dairy products (except yogurt).
- Some vegetables, especially brown beans, cauliflower, brussels sprouts, broccoli, and cabbage.
- Bran.
- Dried fruits.
- Sugar substitutes in sugar-free foods and candies; for example, sorbitol.

A change in eating habits and diet usually takes care of most gas problems. If the condition continues, however, see your doctor to check for other possible causes, such as medication.

This content is reviewed periodically and is subject to change as new health information becomes available. The information is intended to inform and educate and is not a replacement for medical evaluation, advice, diagnosis, or treatment by a health care professional.

Copyright © Clinical Reference Systems 2002

Adult Health Advisor
Senior Health Advisor