1015 S. Hackett Rd.Waterloo, Iowa 50701319.234.5990

CedarValleyGI.com



Ravi Mallavarapu, MD Srinivas Kalala, MD Arun Muthusamy, MD Suhag Patel, MD Moaz Sial, MD Barbara Burkle, ARNP Tracy Elliott, ARNP Kelli DeSerano, ARNP

Colonoscopy Prep Instructions - Osmoprep

(Please also see bowel prep instructions - attached)

- ** There will be instructions on the prescription box that you get from the pharmacy. Please DO NOT use those instructions. Please use the instructions that we provide to you. **
- 7 days prior to the procedure (unless otherwise instructed) stop taking the following:

 Ticlid (ticlopidine) ~ Plavix (clopidogrel) ~ Aggrenox (aspirin/dipyridamole) ~ Effient (prasugrel)
- 5 days prior to the procedure (unless otherwise instructed) stop taking the following:

 Coumadin (warfarin) ~ Iron supplements ~ Metamucil/other fiber supplements ~ Products that contain olestra
- **2 days prior to the procedure** (unless otherwise instructed) stop taking the following: *Pradaxa* (dabigatran etexilate) ~ *Xarelto* (rivaroxaban) ~ *Eliquis* (apixaban)
- ** If you forget to discontinue any of these medications, please call our office between the hours of 8:30 a.m. and 5:00 p.m. or let the nurse know when you arrive for your procedure. It is not necessary to call after normal business hours. **

If you are diabetic:

- Take ½ of your usual dose of medication or insulin in the morning on the day prior to the exam (your prep day).
- Do not take your evening dose of medication or insulin on the day prior to the exam.
- Do not take any medication or insulin on the morning of your exam.
- If you usually do Accu-Chek readings at home, please do one in the morning prior to coming to your appointment on the day of your exam.

2 Days prior to your procedure: Date:
 Take a dose of MiraLax in the evening to ensure a good prep. It should not give you diarrhea or interfere with you daily activities. (You can get Miralax from your local drug store or obtain a complimentary packet from the Cedar Valley Digestive Health Center at 1015 S. Hackett Rd. in Waterloo).
The day prior to your procedure: Date:

- DRINK ONLY CLEAR LIQUIDS THE ENTIRE DAY. Your preparation will work better if you drink extra fluids all day, so
 drink fluids starting first thing in the morning. (Note: if you are on fluid restrictions, please let us know.)
- Drink an 8-ounce glass every hour at minimum. (It is important that you drink plenty of liquids to ensure that the laxative will work.)

Thin	as '	VOII	mav	have:
	yu	you	IIIuy	Huvc.

Apple Juice ~ White grape juice ~ Clear broth ~ Boullion ~ Jello ~ Popsicles ~ Water ~ Black coffee ~ Tea ~ Hi-C ~ Gatorade ~ Lemonade ~ Iced tea ~ Soda ~ Kool-Aid

Things you may not have:

Solid Foods ~ Juices with pulp ~ Alcoholic beverages ~ Milk ~ Milk products ~ Caffeine (Limit 3 per day) ~ Items that are red or purple

Night before your procedure: Date:
Start your laxative at 6:00 pm – see page 3 for further preparation instructions.
Morning of your procedure: Date:
Take the second dose of laxative 4 hours prior to your arrival time (plus drive time) – see page 3 for further preparation instructions. Even if your stools were clear after laxative dose last evening, you <u>must</u> drink the second laxative dose.
Continue to drink clear liquids until Your stools should be clear or yellow, so that you can see through to the bottom of the toilet following a bowel movement. If you are having stools that are brown or contain pieces, call our office at 319-234-5990.

*Take medications 2 hours after drinking your last dose of laxative.

*You may take heart, blood pressure, seizure, anti-rejection, and/or anti-anxiety/depression medication the morning of your procedure. If there are any medications you feel should not be missed, please call the office to check. When you come in, let the nurse know what medications you have taken that day.

- 1. If you use inhalers, please bring them with you.
- 2. You must have a driver due to thesedation you will receive. You will not be able to drive or work until the next day.
- 3. Wear loose, comfortable clothes; you may feel bloated after the procedure.

Date of procedure:	Arrival time
Date of procedure.	Allival tille

** The time that we give you is the ARRIVAL time. The procedure will begin roughly one hour after you arrive. **

** PLAN TO BE HERE HALF THE DAY **

Location: Cedar Valley Digestive Health

1015 S. Hackett Rd. Waterloo, IA 50701

Grundy County Hospital 201 E. J Ave. (Main Entrance #1)

Buchanan County Health Center 1600 1st St. East Independence

Follow up office visit:

If you have any questions or concerns, please call our office at 319-234-5990.

Office hours are Monday–Friday from8:30 a.m.-5:00 p.m., with phones answered 24 hours per day for emergencies.

Or visit our website at: www.myhealthatcvms.com

Please notify your insurance carrier of your upcoming outpatient procedure so you are aware of your benefits.



CedarValleyGI.com