

## Extended Preparation Protocol For Colonoscopy

You should start this 2 days prior to your scheduled colonoscopy. Date: \_\_\_\_\_

These instructions should be used in addition to the colonoscopy prep instructions given to you by the Digestive Health Center nursing staff to ensure optimal preparation for your colonoscopy. This prep is used:

- If you have had poor prep for colonoscopy in the past.
  - If you have a tendency for chronic constipation.
  - If your provider feels you need extra prep prior to your colonoscopy.
- 
- You may have a light breakfast and lunch of soft, easily digestible foods (cooked cereal, eggs, yogurt, cooked fruits or vegetables, cottage cheese, macaroni and cheese, puddings, etc.). You may have clear liquids after lunch, but No solid food until after your colonoscopy.
  - Start extra hydration 2 days prior to your colonoscopy prep. Attempt to drink an 8-ounce glass of water or clear liquid every hour. Please let the office know if you have any fluid restrictions.
  - 5:30 pm: Drink one 10 fluid ounce bottle of magnesium citrate. This is an over the counter laxative.
  - 6-9:00 pm: Continue drinking 8 ounces of clear liquids every hour.
  - 9:30 pm: Take 4 Bisacodyl (Dulcolax) tablets with water.
  - The day prior to your colonoscopy, begin the colonoscopy prep instructions you were given. Date: \_\_\_\_\_

Prep:  Nulytely  Suprep